

Research Article

# To legalize cannabis in Ghana or not to legalize? Reviewing the pharmacological evidence

Kwesi Boadu Mensah<sup>1\*</sup> and Paa Kofi Tawiah Adu-Gyamfi<sup>2</sup>

<sup>1</sup>Department of Pharmacology, Faculty of Pharmacy and Pharmaceutical Sciences, College of Health Science, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

<sup>2</sup>Department of Nursing and Midwifery, Faculty of Health and Allied Sciences Pentecost University College, Accra, Ghana

## More Information

**\*Address for Correspondence:** Kwesi Boadu Mensah, Department of Pharmacology, Faculty of Pharmacy and Pharmaceutical Sciences, College of Health Science, Kwame Nkrumah University of Science and Technology, Kumasi-Ghana, Tel: +233-20-8443173; Email: kbmensah.pharm@knust.edu.gh

**Submitted:** 29 November 2019

**Approved:** 09 December 2019

**Published:** 10 December 2019

**How to cite this article:** Mensah KB, Adu-Gyamfi PKT. To legalize cannabis in Ghana or not to legalize? Reviewing the pharmacological evidence. Arch Pharm Pharma Sci. 2019; 3: 082-088.

**DOI:** dx.doi.org/10.29328/journal.apps.1001018

**ORCID:** orcid.org/0000-0002-9649-2497

**Copyright:** © 2019 Mensah KB, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Keywords:** Medical marijuana; Cannabimimetic; Cannabergics; Sub-Sahara Africa; Cannabinoids



## Summary

**Background:** Although illegal, Ghana has a long history of cannabis use. With changing perceptions, advocacy for legalization has increased globally. This study exams pharmacological evidence on the prospects and challenges of decriminalization and /or legalization of cannabis in Ghana.

**Results:** Cannabis and cannabinoids are a “pharmacological enigma” with unique ability to activate at least 3 of the 4 drug receptor super families. This include; inotropic Transient Receptor Potential Vanilloid 1 (TRPV1), metabotropic Cannabinoid Receptors (CB) and nuclear Peroxisome Proliferator Activator Receptors (PPAR). Cannabinoid receptors also dimerize with other receptors creating distinctly new signaling pathways. Cannabis and cannabinoids show good anti- nociceptive, anti-inflammatory, immunosuppressant anti-emetogenic activity and variable anticonvulsant activity. It can play important role in palliative care, some rare intractable epilepsy, multiple sclerosis, cachexia and Opioid Use Disorder. Cannabis precipitates psychosis in individuals with underlying genetic susceptibility. Chronic cannabis use alter the neurobiology of adolescent brain, predisposing them to amotivational syndrome characterized by depersonalization and inhibited motivation for goal directed behavior. Cannabis is also a “gateway drug”; ushering users to “harder” substances of abuse and reinstating extinguished drug seeking behaviours. The recent tramadol abuse in Ghana may have been precipitated by previous and concurrent cannabis use. Furthermore, Ghana’s cannabis may have a higher propensity to induce detrimental effects because of preferential accumulation the psychotropic delta-9-Tetrahydrocannabinol as a result of the high tropical temperature and humidity.

**Conclusion:** There is not sufficient pharmacological evidence supporting criminalization of medical cannabis in Ghana. However, the same evidence does not support legalization of recreational cannabis.

## Introduction

Contemporary media and scientific literature indicate a global surge in interest in the therapeutic effects and recreational uses of cannabis or marijuana [1]. Consequently, perception on cannabis use is rapidly moving towards liberalization with some industrialized countries decriminalizing or legalizing the use [2]. Compared to decriminalization, legalization provides an economic dimension of regulated cannabis cultivation, supply and taxation system. In such jurisdictions, a multimillion dollar cannabis industry is developing, which provides employment and contributes significantly towards Gross Domestic Product. This has led to rechanneling of

police efforts to more relevant crimes and has reduced the hold of criminals on the trade [3]. Surprisingly, in some cultures, small scale cannabis business are overwhelmingly against legalization for fear of losing decades old monopoly to large co-operations [4].

Recently, Ghana, a major drug transit hub, has also grappled with the issues of legalizing of cannabis. The Ghanaian state is secular but deeply conservative socially. Substance abuse is underreported due to stigmatization. Nonetheless, some limited data suggest that Ghana’s cannabis consumption per capita maybe amongst the top 10% globally. Over 4.5 tonne of cannabis were seized in Ghana in 2017 [5]. Reports on



estimated users range between 8 to 21.5% which is higher than the global average of 3.8% [5,6].

Reasons why cannabis use is popular in Ghana is yet to be determined. Economically, cannabis trade is means of “capital formation” as well as a safety net during cocoa price slump [7,8]. In Ghana, cannabis is relatively cheap compared to other substances of abuse and low cannabis pricing correlates well with abuse [9]. Furthermore, the Ghanaian Boarding Senior High School system is a good breeding group for peer influence to cannabis use. In addition, there is an apparent grey interface between Christianity (the predominant religion in Ghana) and the Rastafarian religion, where cannabis has sacramental uses [2].

This work therefore reviews over five decades of pharmacological research on cannabis and cannabinoids to provide a pharmacological perspective about the prospects and challenges of cannabis use in Ghana.

### What is cannabis or marijuana?

Cannabis or marijuana has many local street names including, wee, ganja, ntampi, indian hemp, popo, obonsam tawa (‘the devil’s tobacco’) etc. It may refer to the whole plant or plant part of the dioecious annual flowering plant, *Cannabis sativa* or its closest variety *Cannabis indica* of family *cannabaceae* (previously *Urticaceae*) or hybrid plant of the varieties [10]. *Cannabis sativa* varieties are taller and have higher phytocannabinoids. Cannabinoids, the main phytochemicals of the plant are in the resins of glandular secretions of the female plant [11]. Typically, dried aerial part is smoked to administer lipophilic phytocannabinoids to the lungs and brain within minutes. Occasionally, cannabis hot infusion (tea), or “Lakka”, a popular alcohol extract of cannabis is used although bioavailability of cannabinoids by oral route can be erratic and unpredictable [12].

History about introduction and use of cannabis in Ghana is not clear. There are suggestion that it may have been introduced in early 20<sup>th</sup> Century by returning World War 11 veterans from Indian and Burma. However, the role of Sierra Leonean immigrants and sailors in the propagation and commercialization of cannabis throughout British West African cannot be overemphasized [13]. Not with standing, cannabis has survived and thrived extensively throughout Ghana especially in cocoa growing areas providing income support when global prices slump [8].

### Cannabinoids

Classically, the terms cannabinoid, cannabimimetic or cannabenergics are synonymous and describe a diverse group of structurally dissimilar families of compounds that bind to cannabinoid receptors (CB<sub>1</sub>/CB<sub>2</sub>) or mimic some of the actions of the major psychoactive substance in *Cannabis sativa*. There are 3 major classes of cannabinoids.

**Phytocannabinoids:** Over 100 chemically distinct C<sub>21</sub> terpenophenolics compounds have been identified and isolated from *Cannabis sativa* [14,15]. Gaoni and Mechoulam (1964) first isolated Delta-9-Tetrahydrocannabinol (THC) as the main phytocannabinoid in the plant responsible for its hallucinogenic activity. Other major phytocannabinoids include cannabigerol, cannabichromene, cannabitrilol and Cannabidiol. Cannabidiol has been shown to antagonize Delta-9-THC’s psychotropic effects [16,17].

**The endocannabinoids system:** The system comprises receptors, endogenous ligands, enzymes that synthesize and degrade signaling molecules [18].

Two endogenous molecules *N*-Arachidonylethanolamine (anandamide) and 2-arachidonoylglycerol (2-AG) have been labelled as endocannabinoid prototypes. Anandamide (“amide of “delight”) is obtained by decarboxylation of phosphatidylserine to phosphatidylethanolamine followed by acetylation [19]. This reaction is carried out by the ubiquitous conjugation enzyme *N*-acetyl transferase. It is then followed by a series of reactions involving Phospholipase A<sub>2</sub>, C and D [20]. Similarly, 2-Arachidonoylglycerol (2-AG) is believed to be a degradation product of glycerophospholipids such as inositol phospholipids and several cellular Phospholipase enzymes [21].

**Synthetic cannabinoids:** These cannabinoids were synthesized to aid in elucidating aspects of the endocannabinoid system. However, they have developed into a distinct pharmacological class use for pharmacotherapy and recreation. The core structure is a bi or tricyclic delta-9-THC compound with a substituted central pyran ring [22]. They possess greater affinity for cannabinoid receptors than delta-9-THC. Popular synthetic cannabinoids such as spice, shatter and K2 have been used as adulterants in herbal preparation and snacks to create mass euphoria and psychosis at public gatherings [22]. Continuous structural modification of the parent molecule has made it a challenge for drug regulatory agencies globally.

### Pharmacology of cannabis/cannabinoids

Pharmacology of cannabis and cannabinoids is complex and it is believed to interact with at least 3 of the four major receptor super families which include inotropic Transient Receptor Potential Vanilloid 1 (TRPV1), the metabotropic Cannabinoids receptors (CB), GPCR 55 & 119 as well as the nuclear Peroxisome Proliferator Activator Receptors (PPAR). Molecular techniques have identified two distinct G-protein coupled cannabinoid receptors [23]. The type 1 cannabinoid receptor (CB<sub>1</sub>) is predominantly distributed in regions of CNS involved in cognition, memory, reward, nociception, and motor coordination as wells some peripheral visceral tissues. The well-known tetrad of cannabinoids effects i.e, hypoactivity, hypothermia, antinociception and catalepsy occur as a result of CB<sub>1</sub> receptors activation. The type 2



receptor (CB<sub>2</sub>) is predominantly found on immune and hematopoietic cells [24]. CB<sub>2</sub> has also been identified in some CNS region and its expression appears to be highly inducible by injury and trauma [25]. Generally, CB receptor activation induces cellular hyperpolarization, decrease cellular cyclic Adenosine Monophosphate levels (cAMP) whilst activating extracellular signal-regulated kinases [26].

However, activations of the classical CB receptors by cannabis does not account for all the myriad of observed pharmacological effects. Cannabinoids bind to Transient Receptor Potential Vanilloid 1 (TRPV1) on small diameter primary afferent fibres leading to reduced sensation to pain. Effects of endocannabinoid on TRPV1 are blocked by capsaicin from chilli but not CB<sub>1</sub> receptor antagonist [27]. Cannabinoids also show strong affinity for binding to Peroxisome Proliferator Activated receptors (PPAR) family of nuclear receptor. These receptors play important role in fat and carbohydrate metabolism, transcription of hepatic enzymes [28]. Moreover, the endocannabinoid prototypes are a biosynthetic product of fatty acid metabolism hence share overlapping physiological effects with products of eicosanoid acid [29].

In addition to activating individual receptors, cannabinoid receptors dimerize with other receptors (either homo or heterodimerization) following concurrent activation creating distinctly new signal transduction pathways. This has been reported cannabinoid CB<sub>1</sub> and dopaminergic D<sub>2</sub> receptors [30].

### Medical potential of cannabis and cannabinoids

**Analgesia/Antinociception:** Cannabis has been used for acute, chronic, inflammatory and neuropathic pain [31]. Infact, antinociception is one of the tetrad of behaviors associated with cannabinoids. CB<sub>1</sub> receptors in the peripheral systems, presynaptic primary afferent C- and A $\delta$ -fiber as well postsynaptic dorsal horn neurons of the spinal cord modulate nociceptive transmission [32]. In addition, CB<sub>1</sub> receptor inhibition of hippocampal neurons calcium channels cause alteration in glutamate release in dorsolateral striatum with arguments anti nociceptive effect [33,34]. Cannabinoids also activate opioidergic pathways in modulating pain [35]. There have been suggestions that a separate metabotropic receptor, GPR55, may be involved in modulating mechanical hyperalgesia. During therapy, cannabinoids synergizes well with other analgesics.

**Anti-Inflammatory and immunosuppression:** Overwhelming evidence suggests that cannabis is effective in diseases with underlying chronic inflammation and immune cell dysfunction such as asthma, rheumatoid arthritis, Crohn's disease. The immune effects is mediated CB<sub>2</sub> receptors located on immune cells; activation of which alter cellular and humoral immunity [36]. Furthermore, the detection of CB<sub>2</sub> receptors in CNS suggest a possible role of endocannabinoids

in neuro protection and neuro inflammation [37]. This may explain why cannabinoids have been very effective in the management of multiple sclerosis [38]. In *in vitro* studies, cannabis shows bidirectional actions on immune cells i.e, low doses enhance survival and proliferation whilst high doses exert inhibitory effects. Secondly, cannabinoids stimulate the immunologic immune system mediated by T helper Type 2 whilst inhibiting the cellular defense immune responses mediated by T helper Type 1 [24]. This is reflected by changes in their respective cytokines levels [39].

**Anti-emetogenic effects:** Nausea and vomiting has been a major drawback to cancer chemotherapy. Conventional anti-emetics such as serotonin 5HT<sub>3</sub> receptors antagonist with corticosteroids have been woefully ineffective at relieving delayed or anticipatory symptoms [40]. A neurokinin NK<sub>1</sub> receptor antagonists was introduced with much greater success. However, neither agents used alone or in combination is effective at suppressing nausea caused by chemotherapy [40].

Cannabis has been employed as anti-emetic, anti-nausea agents over centuries [41]. Cannabis and cannabinoids were putatively accepted as anti-emetic, anti-nausea after patients who actively smoked cannabis showed less susceptible to the emetogenic effect of cancer chemotherapeutic agents. Cannabinoids can protect against 'acute', 'retarded' or 'anticipatory' nausea and vomiting reactions induced by cancer chemotherapy [40,41]. This effect is reversed by CB<sub>1</sub> receptor inhibition implying a possible mediation by CB<sub>1</sub>.

Evidence suggest a probable interaction between serotonergic receptors and cannabinoid receptors [42]. More compelling evidence of this interaction is coexpression CB<sub>1</sub> and 5-HT<sub>3</sub> receptors in regions involved in vomiting such as the area postrema, nucleus of the solitary tract (NTS) and the dorsal motor nucleus of the vagus. Peripherally, cannabinoids also induce GIT relaxation which attenuates vomiting by enteric serotonergic impulses. This is important as mechanistically, vomiting occurs when serotonin activates 5-HT<sub>3</sub> on adjacent enteric parasympathetic afferent nerves to activate chemoreceptor trigger zone [42].

**Cannabis and cancer:** Cannabinoids have been employed in cancer mostly for adjunctive and palliative actions [43]. Epidemiological evidence for an association between cannabis use and decreased incidence of cancer is limited and conflicting [43]. The endocannabinoid system appears to be hyperactive in many cancers and pharmacological manipulation of the receptors usually results in antitumour effects [44]. Cannabinoids show a good degree of selectivity in inducing apoptosis in cancer cells especially in lung, breast, pancreatic, prostate and gliomas [45].

Several apoptotic pathways have been proposed. This include phosphoinositide 3-kinase (PI3K), protein kinase B (Akt), ERK signaling pathways, protein p8, stress-related





transcription factors [44,46]. The anti-tumour activity involve both cannabinoid receptors. The tissue receptor ratio and density may account for variation in response. Consequently, cannabis are procarcinogenic in tissue without cannabinoid receptors or those with low receptor density [47].

**Anticonvulsant:** There are multiple anecdotal references to the effectiveness of cannabis in epilepsy. However, the first documented clinical evidence on efficacy was by O'Shaughnessy in 1848 [48]. Although delta -9- THC exhibits powerful anticonvulsant activity, there are also clinical evidence showing pro-convulsant activity [49]. Despite these contradictions, virtually all the major phytocannabinoids demonstrate efficacy in maximum electroshock in murine but not all show effectiveness in chemoshock models [50]. Some authors have postulated that cannabinoids are effective at controlling spastic seizure states and some rare intractable epilepsy but are virtually useless in petit and grand mal seizures [49,50]. A survey amongst 215 epileptics who actively use cannabis should that only 7.4% reported that cannabis improved their seizure condition [51]. The scientific evidence supporting the use of cannabis for the treatment of epilepsy is at best weak.

**Hyperphagic actions:** According to Abel, 1975, the earliest reference to the hyperphagic actions of cannabis was around AD 300. Subsequently, in the last decades cannabis and cannabinoids have been used to promote weight gain in cachexia, HIV wasting and other hyper metabolic states. There are prospects that cannabinoids will may be beneficial in other nutrition related disorders. The mechanism involve enhancement of orosensory acuity of users resulting in preferential overconsumption of sweet foods especially within the first few days of treatment [52,53]. However, animal studies have not been successful at corroborating the human experience and the magnitude and quantum increase in food consumption reported in animal studies were a fraction of the human experience [53].

Concrete scientific evidence implicate CB<sub>1</sub> receptors in brain centers traditionally associated with food such as ventromedial hypothalamus, the nucleustractus solitaries [54,55]. This findings is further re-enforced by the ability of CB<sub>1</sub> specific antagonist to reverse hyperphagia [54]. This effect is stereo specificity with only the L- delta-9-THC being effective [56]. Naloxone blocks cannabis induced hyperphagia; suggestion a possible interaction between opioids and cannabinoids [57].

## Legalisation of cannabis in Ghana–potential challenges

**Neuropsychiatric effects:** A major setback to legalization is cannabis neuropsychiatric effects. Cannabis use is high amongst individuals with schizorhphrenia, dysthymia and major depression [58,59]. In many instances, the use of cannabinoids precedes the development of neuropsychiatric disorder. Again, cannabis may precipitate acute transient

psychosis in persons with preexisting neurological conditions or those with a positive family history of schizophrenia [60]. In such individuals, adolescent cannabis use can increase the risk of schizophrenia by as much as six folds before the age of 26 [60]. However, as to whether cannabis cause neuropsychiatry disorders is still up for debate as there are no properly controlled population based [59]. Another major setback to this findings is concurrent usage of other illicit drugs by cannabis users. It is however noteworthy that not all people who use cannabis develop acute transient psychosis and schizorhphrenia. A complex interplay of environmental (socio-economic and socio-demographic factors) and genetic vulnerability (eg defective Cathecol-O-methyltransferase enzyme) contributes to the development of neuropsychiatric disorder.

Notwithstanding, these epidemiological studies may not predict the effects of cannabis in the Ghanaian context. Although unsubstantiated, Ghana Cannabis has been putatively touted as “High Grade” supposedly due to its high delta-9-THC content. It is known that the high tropical temperature and humidity in Ghana favour the accumulation of psychotropic delta -9-THC over the non-psychogenic cannabidiol (CBD) [61]. At the molecular level cannabidiol antagonizes the psychotropic effects of delta -9-THC [62,63]. The ratio of delta -9-THC/CBD in cannabis inversely correlates to the volume of right hippocampus of the chronic user [64]. Therefore, cannabis with high delta -9-THC/ CBD ration is associated with more schizophrenia-like symptoms [65,66]. All these may suggest that Ghana's cannabis may have a different chemical composition and a possible higher propensity to induce psychosis than that usually used by many researchers.

## Cannabis, the reward system and retrograde signaling

According to the dopamine theory of psychosis, excess dopamine in some cortical and mesolimbic centers of the brain causes psychosis [67]. Subsequently, cannabinoids act on CB<sub>1</sub> receptors to activate mesolimbic dopamine system for reward and reinforcing. However, Spiller, et al. [68], postulate that higher dose of cannabinoids attenuate reward and induce aversion by stimulating CB<sub>2</sub> receptors [68]. This may explain the paradoxically effects such as euphoria or dysphoria, anxiolysis or anxiety reported by some cannabis users.

The neurobiological basis for concomitant use of cannabis and other substances of abuse is ability of Delta-9-THC to facilitate the reward of other substances of abuse [69]. The accumulation dopamine in the Nucleus Accumbens, and Ventral Tegmental Area and the rewarding effects of several illicit drugs can be reversed by CB<sub>1</sub> receptor antagonists. These effects were confirmed by the decreased rewarding effects of drugs of abuse by CB<sub>1</sub>-null mice [70]. In support of this, CB<sub>1</sub> agonist can reinstate extinguished drug seeking behaviours causing previous drug addicts to relapse [71].



Endocannabinoids acting on CB<sub>1</sub> receptors, modulate proximal neurochemical transmission. This results in retrograde suppression of neurotransmissions to protect against excess presynaptic activity and modulating mesolimbic dopamine releasing. Delta-9-THC disrupts this retrograde signaling of endocannabinoids explaining many neuropharmacological effects of cannabis [72].

### Cannabis as a “gateway” to hard drugs

Studies in substance abusers showed a sequential and incremental pattern where cannabis ushers users to addictive “hard” drugs [73]. Many users begin with alcohol or tobacco proceed through cannabis to narcotics, hallucinogens, methamphetamine [74,75]. Fergusson, et al. [76], report a dose dependent relationship between the use of cannabis and use of other illicit drugs. Furthermore, the age of first exposure to cannabis is also a major predisposing factor to the likelihood of the individual using hard drugs. Indeed, it is very possible that the recent abuse of tramadol in Ghana may have been precipitated by previous and concurrent cannabis use. This presupposes that addressing cannabis use in Ghana will putatively address the surge in tramadol abuse. Suffice to say that delaying the use of alcohol and tobacco by adolescents can reduce or delay the use of cannabis and subsequently the use of other hard drugs.

### Cannabis induced-apathy

Perhaps, the major drawback to legalizing cannabis is its long-term impact on young and adolescent brains inducing functional and neuroanatomical changes [64,77]. Cannabis use is associated with psychosocial amotivational syndrome characterized by depersonalisation, derealisation and an inhibited motivation for goal directed behavior [78]. There is strong evidence to show that adolescent cannabis abusers exhibit a blunted emotion, apathy, a reduced affect, sense of detachment, difficulty to concentrate on relevant issues, follow routines or successfully master new material [79]. Such adolescents are more likely to exhibit delinquent behaviours, absenteeism, diminished educational achievement and drop out of school [80,81]. Reduced reward sensitivity due to reduced striatal dopamine biosynthesis is largely associated with this cannabis-induced amotivational syndrome [82].

### Conclusion

Cannabis has several medical uses and has not shown to be inherently more toxic than several medicines in current clinical practice. There is very minimal pharmacological justification for criminalizing medicinal cannabis use. However, the pharmacological evidence available now is skewed against legalization of cannabis for recreational because its “gateway effects”, association with amotivational syndrome, re-ignition of previous extinguished drug seeking behavior and recidivism in former addicts.

### References

- Daniulaityte R, Nahhas RW, Wijeratne S, Carlson RG, Lamy FR, et al. “Time for dabs”: Analyzing Twitter data on marijuana concentrates across the US. *Drug Alcohol Depend.* 2015; 155: 307-311. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/26338481>
- Davenport S, Pardo B.: Reviewing The dangerous drugs act amendment in Jamaica goals, implementation, and challenges. *Int J Drug Policy.* 2016 1; 37: 60-69. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/27596698>
- Dragone D, Prarolo G, Vanin P, Zanella G. Crime and the legalization of recreational marijuana. *J Economic Behavior Organization.* 2019; 159: 488-501.
- Chappell K. As Jamaica looks to cash in on cannabis, Rastafarians fear being left out. Edited by Jumana Farouky and Zoe Tabary. Thomson Reuters. 2019.
- World Drug Report United Nations Office on Drugs and Crime. 2019.
- World Drug Report Multi-year archive. United Nations Office on Drugs and Crime. 2011.
- Bernstein, Henry, Ghana’s drug economy: some preliminary data, *Review of African Political Economy*79. 1999; 13–32.
- Akyeampong E. Diaspora and drug trafficking in West Africa: A case study of Ghana. *African Affairs.* 2005; 104: 429-447.
- WHO Management of substance. WHO Geneva. 2019.
- Clarke RC, Watson DP. Botany of natural Cannabis medicines. Cannabis and cannabinoids: pharmacology, toxicology and therapeutic potential. 2002; 3-13.
- Hillig KW, Mahlberg PG. A chemotaxonomic analysis of cannabinoid variation in Cannabis (Cannabaceae). *Am J Bot.* 2004; 91: 966-975. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/21653452>
- Akyeampong E. What is in a drink? Class struggle, popular culture and the politics of akpeteshie (local gin) in Ghana, 1930–1967. *J African History.* 1996; 37: 215-236.
- Borrofica A. Mental illness and Indian hemp in Lagos, Nigeria, East. *Afr Med J.* 1966; 43: 377–384. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/5965862>
- Hill AJ, Williams CM, Whalley BJ, Stephens GJ. Phytocannabinoids as novel therapeutic agents in CNS disorders. *Pharmacol Ther.* 2012; 133: 79-97. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/21924288>
- Hanus L, Meyer SM, Muñoz E, Tagliatalata-Scafati O, Appendino G. Phytocannabinoids: a unified critical inventory. *Nat Prod Rep.* 2016; 33: 1357-1392. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/27722705>
- Gaoni Y, Mechoulam R. Isolation, structure and partial synthesis of an active constituent of hashish. *J Am Chem Soc.* 1964; 86: 1646-1647.
- Mc Partland JM, Duncan M, Di Marzo V, Pertwee RG. Are cannabidiol and Δ9-tetrahydrocannabinol negative modulators of the endocannabinoid system? A systematic review. *Br J Pharmacol.* 2015; 172: 737-753. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/25257544>
- Puffenbarger RA. Molecular biology of the enzymes that degrade endocannabinoids. *Current Drug Targets-CNS & Neurological Disorders.* 2005; 4: 625-631.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, et al. Isolation and structure of a brain constituent that binds to the



- cannabinoid receptor. *Science*. 1992; 258: 1946-1949.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/1470919>
20. Liu J, Wang L, Harvey-White J, Huang BX, Kim HY. Multiple pathways involved in the biosynthesis of anandamide. *Neuropharmacology*. 2008; 54: 1-7.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17631919>
  21. Sugiura T, Kobayashi Y, Oka S, Waku K. Biosynthesis and degradation of anandamide and 2-arachidonoylglycerol and their possible physiological significance. *Prostaglandins Leukot Essent Fatty Acids*. 2002; 66: 173-192.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/12052034>
  22. Wiley JL, Marusich JA, Huffman JW. Moving around the molecule: relationship between chemical structure and in vivo activity of synthetic cannabinoids. *Life Sci*. 2014; 97: 55-63.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/24071522>
  23. Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature*. 1990; 346: 561-564.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/2165569>
  24. Tanasescu R, Constantinescu CS. Cannabinoids and the immune system: an overview. *Immunobiology*. 2010; 215: 588-597.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/20153077>
  25. Van Sickle MD, Duncan M, Kingsley PJ, Mouihate A, Urbani P, et al. Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science*. 2005; 310: 329-332.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16224028>
  26. Howlett AC. Cannabinoid inhibition of adenylate cyclase: relative activity of constituents and metabolites of marijuana. *Neuropharmacology*. 1987; 26: 507-512.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/3601007>
  27. De Petrocellis L, Orlando P, Moriello AS, Aviello G, Stott C, et al. Cannabinoid actions at TRPV channels: effects on TRPV3 and TRPV4 and their potential relevance to gastrointestinal inflammation. *Acta Physiol (Oxf)*. 2012; 204: 255-266.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/21726418>
  28. Burstein S. PPAR- $\gamma$ : a nuclear receptor with affinity for cannabinoids. *Life Sci*. 2005; 77: 1674-1684.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16005906>
  29. Fimiani C, Liberty T, Aquirre AJ, Amin I, Ali N, et al. Opiate, cannabinoid, and eicosanoid signaling converges on common intracellular pathways nitric oxide coupling. *Prostaglandins Other Lipid Mediat*. 1999; 57: 23-34.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/10367294>
  30. Kearn CS, Blake-Palmer K, Daniel E, Mackie K, Glass M. Concurrent stimulation of cannabinoid CB1 and dopamine D2 receptors enhances heterodimer formation: a mechanism for receptor cross-talk? *Mol Pharmacol*. 2005; 67: 1697-1704.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/15710746>
  31. Turcotte D, Dorze JA, Esfahani F, Frost E, Gomori A, et al. Examining the roles of cannabinoids in pain and other therapeutic indications: a review. *Expert Opin Pharmacother*. 2010; 11: 17-31.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/20001426>
  32. Agarwal N, Pacher P, Tegeder I, Amaya F, Constantin CE, et al. Cannabinoids mediate analgesia largely via peripheral type 1 cannabinoid receptors in nociceptors. *Nat Neurosci*. 2007; 10: 870-879.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17558404>
  33. Twitchell W, Brown S, Mackie K. Cannabinoids inhibit N- and P/Q-type calcium channels in cultured rat hippocampal neurons. *Journal of neurophysiology*. 1997; 78: 43-50.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/9242259>
  34. Gerdeman G, Lovinger DM. CB1 cannabinoid receptor inhibits synaptic release of glutamate in rat dorsolateral striatum. *J Neurophysiol*. 2001; 85: 468-471.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11152748>
  35. Ottani A, Leone S, Sandrini M, Ferrari A, Bertolini A. The analgesic activity of paracetamol is prevented by the blockade of cannabinoid CB1 receptors. *Eur J Pharmacol*. 2006; 531: 280-281.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16438952>
  36. Ashton JC. Cannabinoids for the treatment of inflammation. *Curr Opin Investig Drugs*. 2007; 8: 373-384.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17520866>
  37. Bisogno T, Di Marzo V. Cannabinoid receptors and endocannabinoids: role in neuroinflammatory and neurodegenerative disorders. *CNS Neurol Disord Drug Targets*. 2010; 9: 564-573.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/20632970>
  38. Zajicek JP, Apostu VI. Role of cannabinoids in multiple sclerosis. *CNS Drugs*. 2011; 25: 187-201.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/21323391>
  39. Croxford JL, Yamamura T. Cannabinoids and the immune system: potential for the treatment of inflammatory diseases?. *J Neuroimmunol*. 2005; 166: 3-18.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16023222>
  40. Slatkin NE. Cannabinoids in the treatment of chemotherapy-induced nausea and vomiting: beyond prevention of acute emesis. *J Support Oncol*. 2007; 5: 1-9.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17566383>
  41. Grinspoon L, Bakalar JB. Marijuana as medicine: a plea for reconsideration. *JAMA*. 1995; 273: 1875-1876.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/7776506>
  42. Bolognini D, Rock EM, Cluny NL, Cascio MG, Limebeer CL, et al. Cannabidiolic acid prevents vomiting in *Suncus murinus* and nausea-induced behaviour in rats by enhancing 5-HT1A receptor activation. *Br J Pharmacol*. 2013; 168: 1456-1470.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/23121618>
  43. Sarfaraz S, Adhami VM, Syed DN, Afaq F, Mukhtar H. Cannabinoids for cancer treatment: progress and promise. *Cancer Res*. 2008; 68: 339-342.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/18199524>
  44. Zhu L X, et al.  $\Delta^9$ -tetrahydrocannabinol inhibits antitumor immunity by a CB2 receptor-mediated, cytokine-dependent pathway. *J. Immunol*. 2000; 165: 373-380.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/10861074>
  45. Velasco R, Zharkikh A, Troggio M, Cartwright DA, Cestaro A, et al. A high quality draft consensus sequence of the genome of a heterozygous grapevine variety. *PLoS One*. 2007; 2: 1326.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/18094749>
  46. Carracedo A, Lorente M, Egia A, Blázquez C, García S, et al. The stress-regulated protein p8 mediates cannabinoid-induced apoptosis of tumor cells. *Cancer Cell*. 2006; 9: 301-312.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16616335>
  47. McKallip RJ, Nagarkatti M, Nagarkatti PS.  $\Delta^9$ -tetrahydrocannabinol enhances breast cancer growth and metastasis by suppression of the antitumor immune response. *J. Immunol*. 2005; 174: 3281-3289.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/15749859>
  48. O'Shaughnessy WB. On the preparation of Indian hemp organjah. *Trans Med Physiol Soc Bengal*. 1842; 421-461.
  49. Katona I. Cannabis and endocannabinoid signaling in epilepsy. *Handb Exp Pharmacol*. 2015; 231: 285-316.  
**PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/26408165>





50. Chesher GB, Jackson DM. Anticonvulsant effects of cannabinoids in mice: drug interactions within cannabinoids and cannabinoid interactions with phenytoin. *Psychopharmacologia*. 1974; 37: 255-264. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/4850601>
51. Gordon E, Devinsky O. Alcohol and marijuana: effects on epilepsy and use by patients with epilepsy. *Epilepsia*. 2001; 42: 1266-1272. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11737161>
52. Abel EL. Cannabis: effects on hunger and thirst. *Behav Biol*. 1975; 15: 255-281. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/1106391>
53. Kirkham TC. Cannabinoids and appetite: food craving and food pleasure. *Int Rev Psychiatry*. 2009; 21: 163-171. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/19367510>
54. Jamshidi N, Taylor DA. Anandamide administration into the ventromedial hypothalamus stimulates appetite in rats. *Br J Pharmacol*. 2001; 134: 1151-1154. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11704633>
55. DiPatrizio NV, Simansky KJ. Activating parabrachial cannabinoid CB1 receptors selectively stimulates feeding of palatable foods in rats. *J Neurosci*. 2008; 28: 9702-9709. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/18815256>
56. McLaughlin CL, Baile CA, Bender PE. Cannabinols and feeding in sheep. *Psychopharmacology (Berl)*. 1979; 64: 321-323. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/116274>
57. Williams CM, Kirkham TC. Reversal of  $\Delta^9$ -THC hyperphagia by SR141716 and naloxone but not dexfenfluramine. *Pharmacol Biochem Behav*. 2002; 71: 333-340. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11812541>
58. Degenhardt L, Hall W. Is cannabis use a contributory cause of psychosis? *Can J Psychiatry*. 2006; 51: 556-565. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17007222>
59. Degenhardt L, Hall W, Lynskey M. Testing hypotheses about the relationship between cannabis use and psychosis. *Drug Alcohol Depend*. 2003; 71: 37-48. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/12821204>
60. McGuire PK, Jones P, Harvey I, Williams M, McGuffin P, et al. Morbid risk of schizophrenia for relatives of patients with cannabis-associated psychosis. *Schizophr Res*. 1995; 15: 277-281. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/7632625>
61. Grlic L. A combined spectrophotometric differentiation of samples of cannabis. *Bull Narcot*. 1968; 20: 25-29.
62. Pertwee R.G. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol*. 2008; 153: 199-215. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17828291>
63. Bhattacharyya S, Morrison PD, Fusar-Poli P, Martin-Santos R, Borgwardt S, et al. Opposite effects of delta-9 tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology*. 2010; 35: 764-774. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/19924114>
64. Demirakca T, Sartorius A, Ende G, Meyer N, Welzel H, et al. Diminished gray matter in the hippocampus of cannabis users: possible protective effects of cannabidiol. *Drug Alcohol Depend*. 2011; 114: 242-245. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/21050680>
65. Morgan CJ, Curran HV. Effects of cannabidiol on schizophrenia-like symptoms in people who use cannabis. *Br J Psychiatry*. 2008; 192: 306-307. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/18378995>
66. Di Forti M, Morgan C, Dazzan P, Pariante C, Mondelli V, et al. High-potency cannabis and the risk of psychosis. *Br J Psychiatry*. 2009; 195: 488-491. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/19949195>
67. Van Rossum JM. The significance of dopamine-receptor blockade for the mechanism of action of neuroleptic drugs. *Arch Int Pharmacodyn Ther*. 1966; 160: 492. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/5954044>
68. Spiller KJ, Bi GH, He Y, Galaj E, Gardner EL, et al. Cannabinoid CB1 and CB2 receptor mechanisms underlie cannabis reward and aversion in rats. *Br J Pharmacol*. 2019; 176: 1268-1281. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/30767215>
69. Wenger T, Moldrich G, Furst S. Neuromorphological background of cannabis addiction. *Brain research bulletin*. 2003; 61: 125-128.
70. Covey DP, Wenzel JM, Cheer JF. Cannabinoid modulation of drug reward and the implications of marijuana legalization. *Brain Res*. 2015; 1628: 233-243. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/25463025>
71. Fattore L, Fadda P, Fratta W. Endocannabinoid regulation of relapse mechanisms. *Pharmacol Res*. 2007; 56: 418-427. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/17936008>
72. Wilson RI, Nicoll RA. Endocannabinoid signaling in the brain. *Science*. 2002; 296: 678-682. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11976437>
73. Morral AR, McCaffrey DF, Paddock SM. Reassessing the marijuana gateway effect. *Addiction*. 2002; 97: 1493-1504. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/12472629>
74. Kandel DB, ed. Stages and pathways of drug involvement: examining the gateway hypothesis. 2002.
75. DuPont RL. Getting tough on gateway drugs: a guide for the family. 1984.
76. Fergusson DM, Boden JM, Horwood LJ. Cannabis use and other illicit drug use: testing the cannabis gateway hypothesis. *Addiction*. 2006; 101: 556-569. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/16548935>
77. Shollenbarger SG, Price J, Wieser J, Lisdahl K. Poorer frontolimbic white matter integrity is associated with chronic cannabis use, FAAH genotype, and increased depressive and apathy symptoms in adolescents and young adults. *NeuroImage: Clinical*. 2015; 8: 117-125.
78. Tennant F S, Groesbeck C J. Psychiatric effects of hashish. *Archives of General Psychiatry*, 1972; 27: 133-136.
79. McGlothlin WH, West LJ. The marijuana problem: an overview. *Am J Psychiatry*. 1968; 125: 370-378. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/5667203>
80. Horwood LJ, Fergusson DM, Hayatbakhsh MR, Najman JM, Coffey C, et al. Cannabis use and educational achievement: findings from three Australasian cohort studies. *Drug Alcohol Depend*. 2010; 110: 247-253. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/20456872>
81. Lynskey M, Hall W. The effects of adolescent cannabis use on educational attainment: a review. *Addiction*. 2000; 95: 1621-1630. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/11219366>
82. Bloomfield MA, Morgan CJ, Kapur S, Curran HV, Howes OD. The link between dopamine function and apathy in cannabis users: an [18 F]-DOPA PET imaging study. *Psychopharmacology (Berl)*. 2014; 231: 2251-2259. **PubMed:** <https://www.ncbi.nlm.nih.gov/pubmed/24696078>